

Supplemental Table 3. Details of systemic steroid administration according to each PICS groups

Variable	<b>Total</b> N=18	PICS N=12	Non-PICS N=6	Patients with 3 domains N=5	Patients without 3 domains N=13	
All steroid treatment	<b>15 (83)</b>	12 (100)	3 (50)	5 (100)	10 (77)	
High-dose*	<b>11 (61)</b>	8 (67)	3 (50)	3 (60)	8 (62)	
Low-dose <sup>†</sup>	<b>4 (22)</b>	4 (33)	0 (0)	2 (40)	2 (15)	
None	<b>3 (17)</b>	0 (0)	3 (50)	0 (0)	3 (23)	

  

Variable	Physical- PICS N=10	Non-physical PICS N=8	Mental-PICS N=9	Non-mental PICS N=9	Cognitive PICS N=6	Non-cognitive PICS N=12
All steroid treatment	10 (100)	5 (63)	9 (100)	6 (67)	6 (100)	9 (75)
High-dose*	6 (60)	5 (63)	7 (78)	4 (44)	4 (67)	7 (58)
Low-dose <sup>†</sup>	4 (40)	0 (0)	2 (22)	2 (22)	2 (33)	2 (17)
None	0 (0)	3 (38)	0 (0)	3 (33)	0 (0)	3 (25)

Data are expressed as numbers (percentage)

\* High-dose was pulse therapy of 500–1000 mg methylprednisolone per day for 3 consecutive days.

<sup>†</sup>Low-dose therapy was 1–2 mg/kg of methylprednisolone for 5–7 days.